

# A selection of ongoing work

## Floral evolution

O'Meara\*, Stacey Smith\*, Armbruster,  
Harder, Hardy, Hileman, Hufford, Litt,  
Magallon, Stephen Smith, Stevens,  
Fenster, Diggle

## Flexible trait evolution

O'Meara and Banbury<sup>P</sup>

## Phylogeographic models

O'Meara, Jackson<sup>P</sup>, Morales-Garcia,  
Carstens

\* = equal lead authors

<sup>P</sup> = postdoc mentee

# A selection of ongoing work

## Floral evolution

O'Meara\*, Stacey Smith\*, Armbruster,  
Harder, Hardy, Hileman, Hufford, Litt,  
Magallon, Stephen Smith, Stevens,  
Fenster, Diggle

## Flexible trait evolution

O'Meara and Banbury<sup>P</sup>

## Phylogeographic models

O'Meara, Jackson<sup>P</sup>, Morales-Garcia,  
Carstens

\* = equal lead authors

<sup>P</sup> = postdoc mentee



NPS




TABLE 1. *List of the eight characters used most frequently in the classification of the families of angiosperms*

Primitive	Character	Advanced	Number in Chart
Calyx and corolla both present (dichlamydous).	Corolla or entire perianth absent (mono- or achlamydous).		1
Petals separate (polypetalous).	Petals (sepals in monochlamydous forms) united (sympetalous).		2
Calyx and corolla regular (actinomorphic).	Perianth irregular (zygomorphic).		3
Stamens more numerous than perianth members (polystemonous).	Stamens the same number as or fewer than perianth members (oligostemonous).		4
Carpels separate (apocarpous).	Carpels united (syncarpous).		5
Seeds more than one per carpel (many).	Seeds not more than one per carpel (few).		6
Placentation axial.	Placentation parietal, basal, or free central.		7
Carpels not united with the receptacle or perianth (hypogyny or perigyny).	Carpels united with receptacle, perianth, or both (epigyny).		8

12 families have presumed ancestral state for all eight traits



36 families few stamens (4), fused carpels (5), and one seed per carpel (6)

CHARACTERS	0	4	6	7	1	46	47	14	67	16	17	467	146	147	167	1467
0	12	5	9	1	2	8	1	1				10				
5	20	20	10	9	1	36	19	3		2	3	7	8	3		24
2	1	4						1				1				
3	1	2					2									
8																
25		17	2	1		12	5	1		2	5	3	1		4	
35	1	5	3		8	4				2	1	1				
58	7	10	1	5	1	8	5			3	1	1		5		
23							1			1						
28																
38																
235		4				8	6				1					
258		10	1		1	8	3	1		1	1	4	3		6	
358		7			2	1				1						
238																
2358		5			3	4	1			3						

No families have presumed derived state for all eight traits

Stebbins (1951): Families are clumped in trait space, therefore “**combinations of characteristics** found in the larger, more successful families of flowering plants **evolved through the guidance of natural selection** because they represented particularly successful methods of solving problems of fertilization, seed dispersal or both.”

Stebbins (1951): Families are clumped in trait space, therefore “**combinations of characteristics** found in the larger, more successful families of flowering plants **evolved through the guidance of natural selection** because they represented particularly successful methods of solving problems of fertilization, seed dispersal or both.”

## Continuing questions

- What is the mechanism? Natural selection for particular combinations? Differential diversification?
- Which traits matter most? Petal presence seems like it would be more important than position of the ovule inside a carpel.
- Is the pattern truly non-random?

TABLE 1. *List of the eight characters used most frequently in the classification of the families of angiosperms*

Primitive	Character	Advanced	Number in Chart
Calyx and corolla both present (dichlamydous).	Corolla or entire perianth absent (mono- or achlamydous).		1
Petals separate (polypetalous).	Petals (sepals in monochlamydous forms) united (sympetalous).		2
Calyx and corolla regular (actinomorphic).	Perianth irregular (zygomorphic).		3
Stamens more numerous than perianth members (polystemonous).	Stamens the same number as or fewer than perianth members (oligostemonous).		4
Carpels separate (apocarpous).	Carpels united (syncarpous).		5
Seeds more than one per carpel (many).	Seeds not more than one per carpel (few).		6
Placentation axial.	Placentation parietal, basal, or free central.		7
Carpels not united with the receptacle or perianth (hypogyny or perigyny).	Carpels united with receptacle, perianth, or both (epigyny).		8

**TABLE 1.** *List of the eight characters used most frequently in the classification of the families of angiosperms*

<b>Primitive</b>	<b>Character</b>	<b>Advanced</b>
Calyx and corolla both present (dichlamydeous).	Corolla present	Corolla or entire perianth absent (mono- or achlamydeous).
Petals separate (diheydrous).	Perianth separate	Petals (separate in apetalous forms) united (sympetalous).
Calyx and corolla both alike (homomorphic).	Radial symmetry	Bilateral symmetry
Stamens more numerous than perianth members (polystemonous).	Many stamens	Few stamens
Carpels separate (apocarpous).	Carpels unfused	Carpels united (synergous).
Seeds more than one per carpel (many).		Seeds not more than one per carpel (few).
Placentation axial.		Placentation parietal, basal, or free central.
Carpels not united with the receptacle or perianth (hypogyny or perigyny).	Ovary superior	Carpels united with receptacle, perianth, or both (epigyny).
		Ovary inferior

Corolla present/absent



Perianth unfused/fused



Radial/bilateral symmetry



Many/few stamens





Carpel unfused/fused





Ovary superior/inferior























# All the factors that lead to contemporary distributions?




# All the factors that lead to contemporary distributions?




# All the factors that lead to contemporary distributions?




Two binary traits: 8 transitions, 4 birth, 4 death




Three binary traits: 12 transitions, 8 birth, 8 death



# Four binary traits: 64 transitions, 16 birth, 16 death




Six binary traits: 384 transitions, 64 birth, 64 death




Six binary traits: 384 transitions, 64 birth, 64 death

Care about combinations, but can't estimate full model



# Six binary traits: 384 transitions, 64 birth, 64 death

## Solution: focal areas



# Six binary traits: 384 transitions, 64 birth, 64 death

## Solution: focal areas




# Six binary traits: 384 transitions, 64 birth, 64 death

## Solution: focal areas



# Six binary traits: 384 transitions, 64 birth, 64 death

## Solution: focal areas



Six binary traits: 384 transitions, 64 birth, 64 death  
Solution: focal areas

## Corolla present/absent


## Perianth unfused/fused

## Radial/bilateral symmetry

Many/few stamens

### Carpel unfused/fused

## Ovary superior/inferior



Six binary traits: 384 transitions, 64 birth, 64 death  
Solution: focal areas

## Corolla present/absent


## Perianth unfused/fused

Radial/bilateral symmetry

Many/few stamens

### Carpel unfused/fused

## Ovary superior/inferior



Six binary traits: 384 transitions, 64 birth, 64 death  
Solution: focal areas

Corolla present/absent


## Perianth unfused/fused

## Radial/bilateral symmetry



Many/few stamens

### Carpel unfused/fused

### Ovary superior/inferior




Six binary traits: *4 transitions, 2 birth, 2 death*  
Solution: focal areas




# Variety of ways to assemble focal areas

(remember, just ways to group rates, so all models remain comparable)


Single character: 0••••



Four characters: 0•11•0



Pair of characters: ••11••



# FaMuSSE: focal area MuSSE


- 5 transition models x 6 diversification models x  $3^6$  focal areas, minus models without enough data
- Total of 19,657 models evaluated
- Estimate likelihood using diversitree (Fitzjohn, 2012) and specified model
- Calculate AIC & Akaike weight

# FaMuSSE: focal area MuSSE

- 5 transition models x 6 diversification models x  $3^6$  focal areas, minus models without enough data
- Total of 19,657 models evaluated
- Estimate likelihood using diversitree (Fitzjohn, 2012) and specified model
- Calculate AIC & Akaike weight

Akaike Weight	Cumulative Akaike Weight	Focal area	Diversification parameters	Transition parameters
0.274	0.274	Corolla present, bilateral symmetry, stamens few	4	1
0.245	0.519	Bilateral symmetry	4	4
0.202	0.721	Bilateral symmetry	4	1
0.128	0.849	Bilateral symmetry, stamens few	4	1
0.090	0.939	Corolla present, bilateral symmetry	4	1
0.024	0.963	Bilateral symmetry	4	2
0.014	0.977	Corolla present, bilateral symmetry, stamens few	4	2
0.009	0.985	Bilateral symmetry	3	4
0.003	0.988	Corolla present, bilateral symmetry	4	2
0.003	0.991	Corolla present, bilateral symmetry, stamens few	4	4


Model averaged results, looking at three characters that matter  
(corolla presence, bilateral symmetry, few stamens)





## Summary:


- Bilateral symmetry important for floral diversification, but so is having few stamens and having a corolla: character combinations matter
- All of these relate to precision of pollen placement: perhaps enable faster speciation?
- Differential diversification rate (“species sorting”) more important than differential transition rates (intraspecific natural selection, mutation, or drift)

# All the factors that lead to contemporary distributions?













Root combination (corolla present, radial symmetry, many stamens)



Fast combination (corolla present, bilateral symmetry, few stamens)







## Summary:

- Bilateral symmetry important for floral diversification, but so is having few stamens and having a corolla: character combinations matter
- All of these relate to precision of pollen placement: perhaps enable faster speciation?
- Differential diversification rate (“species sorting”) more important than differential transition rates (intraspecific natural selection, mutation, or drift)
- Flowers aren’t all at fastest diversifying combination because it took millions of years to evolve that
- Macroevolution limited by microevolutionary variation

## Relevance:

- Go after important questions (macro vs micro, repeatability, non-equilibrium, etc.)

## Relevance:

- Go after important questions (macro vs micro, repeatability, non-equilibrium, etc.)
- Don't let lack of approaches stop progress: build new ones
  - Dating large trees (Smith & O'Meara, treepl, 2012)
  - Multilevel random sampling of taxonomy and GenBank
  - Multiple character \*SSE
  - Forward in time simulations for millions of species

## Relevance:

- Go after important questions (macro vs micro, repeatability, non-equilibrium, etc.)
- Don't let lack of approaches stop progress: build new ones
  - Dating large trees (Smith & O'Meara, treepl, 2012)
  - Multilevel random sampling of taxonomy and GenBank
  - Multiple character \*SSE
  - Forward in time simulations for millions of species
- Answer important questions