


Nicole M. Foley, Richard G. Rasulis, Zoya Wani, Mayra N. Mendoza Cerna, Henrique V. Figueiró, Klaus Peter Koepfli, Terje Raudsepp & William J. Murphy. 2025. "An ancient recombination desert is a speciation supergene in placental mammals" *Nature* https://doi.org/10.1038/s41586-025-09740-2

Class focus area: Genomics and a speciation desert

EEB603: Brian O'Meara

All quotes and images from the above paper unless otherwise noted



**Figure 1. Chromosome connections in meiosis.** Kinetochores attach homologous chromosomes to opposite halves of the spindle. Homologs are held together by chiasmata, in which recombinant chromatids cross each other. Sisters are held together by cohesins and possibly by catenation of centromeric DNA threads, which have been observed in human mitosis. Cohesion is released in two steps: on chromosome arms to resolve chiasmata and separate homologs in the first meiotic division; and around centromeres to separate sisters in the second meiotic division. doi:10.1371/journal.pbio.1000326.g001

## Talbert & Henikoff (2010)







Fig. 1. A cartoon depicting a typical workflow using ReLERNN's four modules (shaded boxes) for (A) individually sequenced genomes or (B) pooled sequences. ReLERNN can optionally (dotted lines) utilize output from stairwayplot, SMC++, and MSMC to simulate under a demographic history with msprime. Training inlays show the network architectures used, with the GRU inlay in (B) depicting the gated connections within each hidden unit. Here, r, z,  $h_{t}$ , and  $\tilde{h}_{t}$  are the reset gate, update gate, activation, and candidate activation, respectively (Cho et al. 2014). The genotype matrix encodes alleles as reference (-1), alternative (1), or padded/missing data (0; not shown). Variant positions are encoded along the real number line (0-1).



Wikipedia gamma distribution








Fig. 1. Discrete approximation to the gamma distribution  $G(\alpha,\beta)$ , with  $\alpha = \beta = \frac{1}{2}$ . Four categories are used to approximate the continuous distribution, with equal probability for each category. The three boundaries are 0.1015, 0.4549, and 1.3233, which are the percentage points corresponding to  $p = \frac{1}{4}$ ,  $\frac{2}{4}$ ,  $\frac{3}{4}$ . The means of the four categories are 0.0334, 0.2519, 0.8203, 2.8944. The medians are 0.0247, 0.2389, 0.7870, 2.3535, and these are scaled to get 0.0291, 0.2807, 0.9248, and 2.7654, so that the mean of the discrete distribution is one.


Yang 1994 "Maximum Likelihood Phylogenetic Estimation from DNA Sequences with Variable Rates over Sites: Approximate Methods" J Molecular Evolution 39:306-314

















- Deep learning algorithm?
- Why just X, why not Y?
- Syntenic alignments? Haldane's rule? Chromosome enrichment? Dosage compensation?
- Is there evidence this occurs beyond mammals?
- Also how uncertain are we about some of the realtionships within mammals? I know the broad (Afrotheria, Laurasiathera, etc.) seem stable from what I have seen, but how much uncertanty do we have?
- It makes me wonder which region is so well-conserved in plants (chloroplast?), given that they don't exhibit much sexual isolation